



E1878
JACC March 27, 2012
Volume 59, Issue 13



Quality of Care and Outcomes Assessment

EXERCISE CAPACITY AND CARDIAC FUNCTION OF HEALTHY OBESITY ARE AS GOOD AS THOSE OF NON-OBESE PERSONS

ACC Moderated Poster Contributions
McCormick Place South, Hall A
Monday, March 26, 2012, 11:00 a.m.-Noon

Session Title: Predicting Quality Outcomes
Abstract Category: 31. Quality of Care and Outcomes Assessment
Presentation Number: 1253-151

Authors: *Mi-Seung Shin, Bong ROUNG Kim, Kim Myeonggun, Jeonggeun Moon, Woong Chol Kang, Gachon University Gil Hospital, Incheon, South Korea, Seoul Medical Center, Seoul, South Korea*

Background: Studies including healthy uncomplicated obesity on alteration of cardiac structure and function are rare. The purpose was to evaluate the impact of obesity on cardiac structure and function in preserved exercise capacity.

Methods: To assess the influence of obesity on LV function, total 303 subjects without evidence of heart, lung and kidney disease were evaluated; 147 obese subjects (74 men) defined as a body mass index (BMI) ≥ 25 kg/m² and 129 non-obese controls (78 men). All subjects underwent transthoracic echocardiography and treadmill exercise test. The mitral early and late diastolic peak velocity (E, A), decelerating time (DT), mitral annular early and late diastolic peak velocity (Ea, Aa), LV end-diastolic dimension (LVEDD), LV ejection fraction (LVEF) were measured. Among those, data of 94 men and 74 women with preserved exercise capacity (≥ 12 METs in men, ≥ 11 METs in women) were analyzed.

Results: In overall 303 subjects, obese men and women showed significantly increased LVEDD, LV mass index and blood pressure (BP) but decreased diastolic function and exercise METs compared with those of non-obese men and women. E/Ea ratio showed negative correlation with METs in obese men ($r = -0.32$, $p < 0.001$) and women ($r = -0.40$, $p < 0.001$). Obese men with preserved exercise capacity ($n = 37$) showed increased LVEDD and BP compared to non-obese men with preserved exercise capacity ($n = 57$). However, cardiac systolic and diastolic function and METs were not significantly different. Obese women with preserved exercise capacity ($n = 30$) showed increased BP compared to non-obese women with preserved exercise capacity ($n = 44$). However, cardiac diameter and systolic and diastolic function were not significantly different.

Conclusions: Obesity was associated with LV structural change, diastolic dysfunction and low exercise capacity. Obese persons with preserved exercise capacity have similar cardiac function compared with non-obese persons with preserved exercise capacity. Exercise capacity and cardiac function of healthy obesity are as good as those of non-obese persons.